

# Assessment of upper airways measurements in patients with mandibular skeletal Class II malocclusion

Nayanna Nadja da Silva<sup>1</sup>, Rosa Helena Wanderley Lacerda<sup>2</sup>, Alexandre Wellos Cunha Silva<sup>3</sup>, Tania Braga Ramos<sup>4</sup>

DOI: <http://dx.doi.org/10.1590/2177-6709.20.5.086-093.oar>

**Objective:** Mandibular Class II malocclusions seem to interfere in upper airways measurements. The aim of this study was to assess the upper airways measurements of patients with skeletal Class II malocclusion in order to investigate the association between these measurements and the position and length of the mandible as well as mandibular growth trend, comparing the Class II group with a Class I one.

**Methods:** A total of 80 lateral cephalograms from 80 individuals aged between 10 and 17 years old were assessed. Forty radiographs of Class I malocclusion individuals were matched by age with forty radiographs of individuals with mandibular Class II malocclusion. McNamara Jr., Ricketts, Downs and Jarabak's measurements were used for cephalometric evaluation. Data were submitted to descriptive and inferential statistical analysis by means of SPSS 20.0 statistical package. Student's t-test, Pearson correlation and intraclass correlation coefficient were used. A 95% confidence interval and 5% significance level were adopted to interpret the results.

**Results:** There were differences between groups. Oropharynx and nasopharynx sizes as well as mandibular position and length were found to be reduced in Class II individuals. There was a statistically significant positive correlation between the size of the oropharynx and Xi-Pm, Co-Gn and SNB measurements. In addition, the size of the nasopharynx was found to be correlated with Xi-Pm, Co-Gn, facial depth, SNB, facial axis and FMA.

**Conclusion:** Individuals with mandibular Class II malocclusion were shown to have upper airways measurements diminished. There was a correlation between mandibular length and position and the size of oropharynx and nasopharynx.

**Keywords:** Angle Class II malocclusion. Oropharynx. Nasopharynx. Airway obstruction.

<sup>1</sup>Specialist in Orthodontics, Associação Brasileira de Odontologia (ABO-PB), João Pessoa, Paraíba, Brazil.

<sup>2</sup>Coordinator, Postgraduate Program in Orthodontics, Associação Brasileira de Odontologia (ABO-PB), João Pessoa, Paraíba, Brazil.

<sup>3</sup>Professor, Postgraduate Program in Orthodontics, Associação Brasileira de Odontologia (ABO-PB), João Pessoa, Paraíba, Brazil.

<sup>4</sup>Professor of Orthognathic Surgery, Postgraduate Program in Orthodontics, Associação Brasileira de Odontologia (ABO-PB), João Pessoa, Paraíba, Brazil.

**How to cite this article:** Silva NN, Lacerda RHW, Silva AWC, Ramos TB. Assessment of upper airways measurements in patients with mandibular skeletal Class II malocclusion. *Dental Press J Orthod*. 2015 Sept-Oct;20(5):86-93. DOI: <http://dx.doi.org/10.1590/2177-6709.20.5.086-093.oar>

**Submitted:** January 13, 2015 - **Revised and accepted:** May 18, 2015

» The authors report no commercial, proprietary or financial interest in the products or companies described in this article.

**Contact address:** Rosa Helena Wanderley Lacerda  
Associação Brasileira de Ortodontia  
Rui Barbosa 38, João Pessoa, Paraíba - CEP: 58.020-040 - Brazil  
E-mail: [rhelenawanderley@msn.com](mailto:rhelenawanderley@msn.com)

## INTRODUCTION

Skeletal Class II malocclusion is a dentofacial deformity caused by a growth disorder of the bones frequently associated with mandibular retrusion relative to upper facial structures.<sup>1</sup> This deformity is also associated with functional disorders, mainly affecting upper airways and the temporomandibular joint.<sup>2,3</sup>

Patients with skeletal Class II malocclusion who have this deformity due to deficiency in mandibular growth present with a retrognathic mandible either because of growth vector or by deficient mandibular length.

According to Muto et al,<sup>4</sup> craniofacial abnormalities, including mandibular retrognathism, short mandibular body length and backward/downward rotation, can lead to decreased pharyngeal airway. These findings indicate that nasopharyngeal obstruction may be related to changes in mandibular morphology.<sup>5</sup>

The study of upper airways and their relationship with mandibular position and size is extremely important in orthodontic diagnosis because of their association with obstructive respiratory disorders, especially sleep apnea. This knowledge is definitive to the indication of mandibular advancement, whether orthopedic or surgical, for treatment of these disorders.

Several studies have been carried out with a view to measuring the pharyngeal airway; however, comparison with Class I individuals and the correlation between the variables involved in Class II malocclusion and airways measurements are still scarce, which encouraged the present study.

## MATERIAL AND METHODS

This study was submitted to and approved by the Ethics Committee on Human Research through Plataforma Brasil, following the norms of the law 466/2012, under approval protocol #835.928.

The sample comprised 80 digital lateral cephalograms belonging to 80 patients of both sexes, without associated abnormalities, aged between 10-17 years, with a mean age of 12.3 years, treated by postgraduate orthodontic students (ABO/PB, Brazil). Of the 80 images, 40 were from patients with mandibular Class II malocclusion, whose diagnosis was confirmed by Xi-Pm, Co-Gn, Go-Me, facial depth and SNB measurements (at least three of these measures should be reduced so that the image would not be withdrawn from the sample). The other 40 radiographs belonged to Class I individuals. Groups were matched by age.

Anatomical tracings of all radiographic images were made on acetate paper, in a dark room, by an examiner using graphite pencil (point 0.3). Each film was traced by one investigator and checked by a second one, so as to verify the accuracy of anatomical outline determination and landmark placement. Measurements of mandibular length and spatial position, as well as size of nasopharyngeal and oropharyngeal airways, were taken using the cephalograms (Fig 1, Table 1).

Measurements were taken twice, with a 10-day interval in between, with the aid of a millimeter ruler and a 180° protractor. The first assessment was carried out with the entire sample while the second one was carried out with 30% of the sample.

Procedures of statistical inference were performed based on parametric statistics. Correlation coefficient and intra-class correlation coefficient (ICC) were used to assess intra-examiner agreement. The choice for statistical test was based on normal distribution of data, according to Kolmogorov-Smirnov normality test ( $p > 0.05$ ). Intergroup comparison was performed by Student's t-test and Pearson r correlation coefficient. For descriptive procedures, absolute and relative data and measurements of central tendency and variability were presented. A 95% confidence interval and 5% significance level ( $p < 0.05$ ) were adopted to interpret the results. Data were submitted to SPSS 20.0 statistical package for Windows and analyzed by means of descriptive and inferential statistics.




Figure 1 - Cephalogram and cephalometric measurements used.

**Table 1** - Cephalometric measurements used.

Measure	Clinical standard	Appropriate age	Description
1. Xi-Pm	65 ± 3 mm	9 years (1.6/year)	Axis of the mandibular body – a line extending from point Xi to the mental protuberance.
2. Facial axis (Ba.NA x Frankfurt)	90 ± 3°	Does not change upon growth	Provides the direction of growth of the chin and the ratio between facial height and depth.
3. Facial depth (NA-Pog x Frankfurt)	87 ± 3°	9 years (0.33/ year)	Indicates the anteroposterior position of the mandible.
4. Co-Gn (Effective mandibular length)			Consists in the geometric relationship between the maxillomandibular length, directly linked either to patient's age or sex.
5. Oropharynx	10 to 12 mm		Measured by the width of the pharynx at the point where the posterior border of the tongue (in the radiograph) crosses the lower border of the mandible up to the posterior pharyngeal wall.
6. Nasopharynx	Mixed dentition: 12 mm Permanent dentition: 17.4 mm		It is measured linearly from a midpoint on the posterior wall of the soft palate to the posterior pharyngeal wall where there is the greatest closure of the airway.
7. Ar-Go	44 mm	11 years (male: 1.01 – 7.2) (female: 0.71 – 4.2)	Height of the mandibular ramus.
8. Go-Me	71 mm	11 years (male: 1.11 – 7.11) (female: 0.73 – 3.12)	Length of the mandibular body.
9. SNB	80°		Anteroposterior position of the mandible in relation to the base of the cranium.

## RESULTS

In order to assess the reliability of measurements of oropharynx and nasopharynx, mandibular length, mandibular position and direction of mandibular growth, the examiner conducted two assessments which were followed by determination of intraexaminer agreement. This calculation was done using intraclass correlation coefficient (ICC). Results were statistically significant and indicated intraclass coefficients ranging from 0.97 (facial depth) and 1.00 (oropharynx), thereby denoting strong intraexaminer agreement (Table 2).

As for upper airways measurements, statistically significant differences were found between both groups ( $p < 0.001$ ). That is, the size of nasopharynx and oropharynx is reduced in Class II individuals (Fig 2).

The same results were observed for mandibular length, with significant differences between groups. The following measurements were found to be greater in Class I individuals: Xi-Pm, Co-Gn and Go-Me (Table 3, Fig 3).

As shown in Table 3, measurements of mandibular position also indicated significant differences between

groups, with facial depth and SNB being greater among Class I individuals (Table 3). These results are graphically shown in Figure 4.


Measurements related to the direction of mandibular growth also differed significantly between groups. Facial axis and Ar-Go were greater in Class I individuals, while FMA was found to be greater in Class II individuals (Table 3, Fig 5).

In order to assess the correlation between oropharynx/nasopharynx size and mandibular length, position as well as growth, Pearson  $r$  correlation coefficient was performed.

Significant positive correlations were observed between the oropharynx and Xi-Pm, Co-Gn and SNB. Moreover, there were also correlations between the nasopharynx and Xi-Pm, Co-Gn, facial depth, SNB, facial axis and Ar-Go. Given that such correlations were positive, it is concluded that the greater the measurements of upper airways, the greater the variables, as reported herein. Correlation coefficients ranged from 0.24 to 0.37; thus, indicating weak to moderate correlations between variables (Table 4).

**Table 2** - Assessment of intraexaminer agreement.


Measures	ICC	p	Intpretation
Xi-Pm	0.99	< 0.001	Strong intra-examiner agreement
Co-Gn	0.99	< 0.001	Strong intra-examiner agreement
Go-Me	0.99	< 0.001	Strong intra-examiner agreement
Facial depth	0.97	< 0.001	Strong intra-examiner agreement
SNB	0.99	< 0.001	Strong intra-examiner agreement
Facial axis	0.99	< 0.001	Strong intra-examiner agreement
Ar-Go	0.99	< 0.001	Strong intra-examiner agreement
FMA	0.99	< 0.001	Strong intra-examiner agreement
Oropharynx	1.00	-	Perfect agreement
Nasopharynx	0.99	< 0.001	Strong intra-examiner agreement




**Figure 2** - Assessment of upper airway measurements of Class I and Class II groups.

**Table 3** - Assessment of upper airways measurements, mandibular length, mandibular position and direction of mandibular growth of each group.

General measures	Specific measures	Class I		Class II		t (p)
		Mean ± SD	Min-Max	Mean ± SD	Min-Max	
Upper airways	Oropharynx	12.2±2.5	7 – 18	8.6±1.7	5 – 13	7.4 (< 0.001)
	Nasopharynx	9.4±1.9	6 – 14	6.7±1.9	3 – 12	6.2 (< 0.001)
Mandibular length	Xi-Pm	77.5±5.4	67 – 94	72.4±4.8	61 – 83	4.4 (< 0.001)
	Co-Gn	115.6±6.7	100 – 134	109.7±7.5	94 – 128	3.6 (< 0.001)
	Go-Me	73.5±11.2	13 – 90	69.4±4.9	57 – 80	2.0 (0.04)
Mandibular position	Facial depth	89.4±2.4	84 – 94	86.1±2.5	79 – 91	5.8 (< 0.001)
	SNB	79.7±2.9	74 – 88	74.5±2.9	68 – 84	7.8 (< 0.001)
Direction of the mandibular growth	Facial axis	90.5±3.7	80 – 100	87.4±3.6	78 – 93	3.8 (< 0.001)
	Ar-Go	43.9±4.1	37 – 51	40.5±4.8	31 – 50	3.3 (0.001)
	FMA	24.9±3.8	14 – 31	27.0±4.9	14 – 36	2.0 (0.04)



**Figure 3** - Assessment of mandibular length of Class I and Class II groups.



**Figure 4** - Assessment of mandibular position of Class I and Class II groups.




Figure 5 - Assessment of mandibular growth of Class I and Class II groups.

Table 4 - Correlation between upper airways measurements and mandibular length, position as well as direction of mandibular growth in both groups.

Measures	r	P	%	Interpretation
<b>Oropharynx</b>				
Xi-Pm	0.31	0.004	9.6%	Significant, positive and moderate correlation
Co-Gn	0.24	0.02	5.7%	Significant, positive and weak correlation
Go-Me	0.13	0.23	1.6%	There was no correlation between variables
Facial depth	0.21	0.06	4.4%	There was no correlation between variables
SNB	0.37	0.001	13.6%	Significant, positive and moderate correlation
Facial axis	0.12	0.26	1.4%	There was no correlation between variables
Ar-Go	0.12	0.28	1.4%	There was no correlation between variables
FMA	-0.07	0.52	0.4%	There was no correlation between variables
<b>Nasopharynx</b>				
Xi-Pm	0.37	0.001	13.6%	Significant, positive and moderate correlation
Co-Gn	0.32	0.003	10.2%	Significant, positive and moderate correlation
Go-Me	0.18	0.11	3.2%	There was no correlation between variables
Facial depth	0.29	0.009	8.4%	Significant, positive and weak correlation
SNB	0.34	0.002	11.5%	Significant, positive and moderate correlation
Facial axis	0.28	0.01	7.8%	Significant, positive and weak correlation
Ar-Go	0.29	0.007	8.4%	Significant, positive and weak correlation
FMA	-0.13	0.24	1.6%	There was no correlation between variables

## DISCUSSION

Although some recent studies have reported a need for tridimensional evaluation by magnetic resonance,<sup>6,7,8</sup> its high cost and lack of standardization of patient's head position still hamper the use of this method for research. According to Muto et al,<sup>9</sup> a change of 10° in craniofacial tilt may affect measurement taking in the area of upper airways in approximately 4 mm. Lateral cephalograms have been used in this type of assessment as part of

patients' basic orthodontic records, with the advantage of having low costs and low radiation dose, being of easy access, and providing standardization of measures with high reproducibility for diagnosis.<sup>6,10,11</sup> These advantages render this method common in research,<sup>7,9,12,13,14</sup> which validates the methodology adopted in the present study and allows comparison of results. The reproducibility of the method was confirmed statistically, with strong intraexaminer agreement.

The studied sample comprised patients aged between 10 and 17 years old, with a mean age of 12.3 years, similarly to other studies,<sup>5,12,15,16</sup>. Because there are minor changes in the nasopharynx as a result of growth,<sup>17</sup> the sample was matched by age; thus, avoiding potential bias as regards data interpretation. In terms of sex, groups were similar, although we found three more males than females in the Class II group.

Regarding airways measurements, there were significant differences between groups, with Class I patients having oropharynx and nasopharynx greater in size (Table 3, Fig 2). These findings corroborate the majority of studies found in the literature.<sup>14,18,19,20</sup> The studies by Freitas et al<sup>12</sup> as well as Memon, Fida and Shaikh<sup>21</sup> found no interference of malocclusion in oropharynx and nasopharynx width when they compared Class I to Class II patients. Differences in our results may be related to the methods employed, since those studies included a Class II sample based on dental occlusion and may have included subjects with Class II resulting from maxillary prognathism, whereas in our study, mandibular Class II was confirmed cephalometrically.

In order to have a better understanding of which factors inherent to malocclusion could be related to changes in upper airways, we initially diagnosed differences in skeletal features between groups, as follows: mandibular length (Xi-Pm, Co-Gn and Go-Me), mandibular position (facial depth and SNB), and direction of growth (facial axis, Ar-Go and FMA).

As regards mandibular length, measurements found in the Class I group were greater than those found in the Class II group (Table 3, Fig 3), thereby confirming mandibular Class II diagnosis. These data validate the assumption that mandibular length can be related to the size of upper airways, which is in agreement with Muto et al<sup>4</sup> who pointed out that craniofacial abnormalities, including mandibular retrognathism, short mandibular body and downward rotation, may cause a decrease in the size of airways, as reported by other studies.<sup>9,13,19,22,23</sup> The same behavior was observed in the variables related to spatial position of the mandible. As expected, the mandible in the Class II group was found retropositioned in relation to the cranial base when compared to the Class I group. This information allows us to conclude that both position and length of the mandible, i.e., the effective length of the mandible, must be considered in the diagnosis of patients with

Class II malocclusion. Nevertheless, a greater or less interference of either one of these variables cannot be assumed. In the literature, this comparison is scarce and only cited by a few authors.<sup>1,23,24</sup>

Our study was carried out considering that several others have assessed the association between facial growth pattern and upper airways measurements.<sup>5,12,15,16,19</sup> When comparing Class I and Class II groups, FMA and facial axis indicated an increased vertical trend among Class II individuals as well as a shorter mandibular ramus. According to Jarabak,<sup>25</sup> this finding refers to mandibular morphology with a clockwise growth pattern. This same feature was reported in the study by Joseph et al<sup>15</sup> who used a sample of individuals with Class II malocclusion. This information does not allow us to claim that all mandibular Class II individuals will have a vertical growth trend, although such feature was found in the sample. However, there seem to be an association between vertical pattern and reduced airways measurements, which has already been reported by several studies.<sup>5,12,14,19</sup>

The correlation between oropharynx and nasopharynx was studied separately from other variables, as shown in Table 4. There was a positive correlation between the size of the oropharynx and mandibular length, represented by Xi-Pm and Co-Gn, and the position of the mandible, represented by SNB. In agreement with our findings, studies carried out in the last five years<sup>7,20,23,24</sup> have concluded that mandibular length and position influence airways measurements.

Although Class II malocclusion patients have mostly presented with a vertical growth pattern in relation to Class I individuals, our results could not support a correlation between vertical pattern and a shorter oropharynx. We did not observe a positive correlation between growth pattern measurements (FMA, Ar-Go and facial axis) and the size of the oropharynx, even though there was an association. This is in agreement with the reports by Castro and Vasconcelos.<sup>16</sup> On the other hand, Freitas et al,<sup>12</sup> Zhong et al<sup>19</sup> as well as Ucar and Uysal<sup>5</sup> found a correlation between growth pattern and the size of the oropharynx.

When assessing airways measurements and growth pattern, Joseph et al<sup>15</sup> showed a correspondence between dolichocephalic individuals and shorter airways, particularly the nasopharynx. This is in agreement with our findings, as seen in Table 5 which shows a significant

positive correlation between Ar-Go values and the size of the nasopharynx. In addition, they showed a positive correlation between Xi-Pm, Co-Gn, facial depth, SNB and facial axis; thus, concluding that mandibular length and position are related to the size of the nasopharynx.

Mandibular retrusion is one of the factors that may cause obstructive sleep apnea syndrome (OSA), characterized by a collapse site hindering the passage of air located in the pharynx. A reduction in this region can be the etiology of this syndrome both in children and adults. Characterized by respiratory disorders and nocturnal snoring, OSA may cause psychological and social impairment for the individual.<sup>11,22,23</sup>

As the results of our study suggest that mandibular length and position as well as the direction of growth can influence measurements of pharyngeal airways, we emphasize the importance of mandibular advancement

in growing children through orthopedics by means of functional appliances; and in adults, with surgical advancement in order to promote enlargement of airways for functional and quality of life improvement, as well as decreased morbidity.<sup>8,13,14,26,27</sup>

## CONCLUSION

- » Individuals with mandibular Class II malocclusion were shown to have upper airways measurements reduced when compared to Class I individuals.
- » Mandibular length is related to a decrease in upper airways measurements. Similarly, anteroposterior positioning of the mandible exerts influence on airways measurements.
- » There was a tendency of facial growth pattern with a positive, but weak correlation with the sizes of the nasopharynx, but not with the oropharynx.



## REFERENCES

- McNamara Jr JA. A Components of Class II malocclusion in child 8-10 years age. *Angle Orthod.* 1981;51(3):177-201.
- Alcazar NMPV, Freitas MR, Janson G, Henriques JFC, Freitas KMS. Estudo cefalométrico comparativo dos espaços naso e bucofaringeo nas más oclusões Classe I e Classe II, Divisão 1, sem tratamento ortodôntico, com diferentes padrões de crescimento. *Rev Dental Press Ortod Ortop Facial.* 2004;9(4):68-76.
- Joseph AA, Elbowaa J, Cisneros GJ, Eisigsb A. Cephalometric comparative study of the soft tissue airway dimensions in persons with hyperdivergent and normodivergent facial pattern. *J Oral Maxillofac surg.* 1998;56:135-9.
- Muto T, Yamazaki A, Takeda S. A cephalometric evaluation of the pharyngeal airway space in patients with mandibular retrognathia and prognathia, and normal subjects. *Int J Oral Maxillofac Surg.* 2008;37(3):228-31.
- Ucar FI, Uysal T. Orofacial dimensions in subjects with Class I malocclusion and different growth patterns. *Angle Orthod.* 2011;81(3):460-8.
- Pirilä-Parkkinen K, Löppönen H, Nieminen P, Tolonen U, Pääkkö E, Pirttiniemi P. Validity of upper airway assessment in children A clinical, cephalometric, and MRI study. *Angle Orthod.* 2011;81(3):433-9.
- Kim Y-J, Hong J-S, Hwang Y-I, Park Y-H. Three-dimensional analysis of pharyngeal airway in preadolescent children with different anteroposterior skeletal patterns. *Am J Orthod Dentofacial Orthop.* 2010;137(3):303-11.
- Schutz TCB, Dominguez GC, Pradella-Hallinan M, Cunha TCA, Tufik S. Class II correction improves nocturnal breathing in adolescents. *Angle Orthod.* 2011;81(2):222-28.
- Muto T, Yamazaki A, Takeda S, Kawakami J, Tsuji Y, Shibata T, Mizoguchi I. Relationship between the pharyngeal airway space and craniofacial morphology, taking into account head posture. *Int J Oral Maxillofac Surg.* 2006;35:132-6.
- Major MP, Flores-Mir C, Major PW. Assessment of lateral cephalometric diagnosis of adenoid hypertrophy and posterior upper airway obstruction: a systematic review. *Am J Orthod Dentofacial Orthop.* 2006;130(6):700-8.
- Bittencourt LRA, Haddad FLM. Diagnóstico e abordagem clínica do paciente com distúrbio respiratório do sono. In: Fabro CD, Chaves CM Jr, Tufik S. *A odontologia na medicina do sono.* Maringá: Dental Press; 2012. cap. 6, p. 144-58.
- Freitas MR, Alcazar NM, Janson G, De Freitas KM, Henriques JFC. Upper and lower pharyngeal airways in subjects with Class I and Class II malocclusions and different growth patterns. *Am J Orthod Dentofacial Orthop.* 2006;130(6):742-45.
- Susarla AM, Abramson ZR, Dodson TB, Kaban LB. Cephalometric measurement of upper airway length correlates with the presence and severity of obstructive sleep apnea. *J Oral Maxillofac Surg.* 2010;68:2846-55.
- Restrepo C, Santamaria S, Pela EZ, Tapias A. Oropharyngeal airway dimensions after treatment with functional appliances in Class II retrognathic children. *J Oral Rehabil.* 2011;38(8):588-94.
- Joseph AA, Elbaum J, Cisneros GJ, Eisig SB. A cephalometric comparative study of the soft tissue airway dimensions in persons with hyperdivergent and normodivergent facial patterns. *J Oral Maxillofac Surg.* 1998;56(2):135-9.
- Castro AMA, Vasconcelos MHF. Avaliação da influência do tipo facial nos tamanhos dos espaços aéreos nasofaríngeo e bucofaringeo. *Rev Dental Press Ortod Ortop Facial.* 2008;13(6):43-50.
- McNamara JA Jr. A method of cephalometric evaluation. *Am J Orthod.* 1984;86(6):449-69.
- Mergen DC, Jacobs RM. The size of nasopharynx associated with normalocclusion and Class II malocclusion. *Angle Orthod.* 1970;40(4):342-6.
- Zhong Z, Tang Z, Gao X, Zeng XL. A comparison study of upper airway among different skeletal craniofacial patterns in nonsnoring Chinese children. *Angle Orthod.* 2010;80(2):267-74.
- El H, Palomo JM. Airway volume for different dentofacial skeletal Patterns. *Am J Orthod Dentofacial Orthop.* 2011;139(6):511-21.
- Memon S, Fida M, Shaikh A. Comparison of different craniofacial patterns with pharyngeal widths. *J Coll Physicians Surg Pak.* 2012;22(5):302-6.
- Schwab RJ, Goldbert AN. Upper airway assessment: radiographic and other imaging techniques. *Otolaryngol Clin North Am.* 1998;31(6):931-68.
- Guarim JA. Evaluation of the growth mandibular in a buccal respirator after the treatment with the use of the orthopedical prefabricated apparel. *Rev Paul Odontol.* 2009;32:15-23.
- Kim JS, Kim JK, Hong SC, Cho JH. Changes in the upper airway after counterclockwise maxillomandibular advancement in young Korean women with class II malocclusion deformity. *J Oral Maxillofac Surg.* 2013;71:1603-5.
- Jarabak JR, Fizzel JA. *Technique and treatment with light wire edgewise appliances.* 2nd ed. St. Louis: Mosby; 1972.
- Lye KW. Effect of Orthognathic Surgery on the Posterior Airway Space (PAS). *Ann Acad Med Singapore.* 2008;37(8):677-82.
- Faria AC, Santos AC, Silva-Júnior SN, Xavier SN, Mello-Filho FV. Change in retrolingual airway space after maxillomandibular advancement surgery in OSA patients: evaluation with magnetic resonance imaging. *Rev Bras Cir Craniomaxilofac.* 2011;14 (3):145-8.