

Factors that favor tridimensional and hermetic obturation of root canal system

Jefferson J. C. **MARION**¹

Déric Meschiari **BATISTA**²

Sueli Satomi **MURATA**³

Valdir de **SOUZA**⁴

Roberto **HOLLAND**⁴

ABSTRACT

Objective: This study aims at demonstrating the possibility and importance of filling different types of accessory canals in different teeth. **Methods:** The sample comprised 11 cases in which detailed knowledge of the inner anatomy of root canal systems associated with good use of the crown-down preparation technique — irrigation with 1% sodium hypochlorite and 17% trisodium EDTA for cleaning and removing smear layer — enabled different types of filling cement to drain. **Results:** Endométhasone,

Sealapex™ and Zinc Oxide - Eugenol filling cements proved to be efficient in root canal filling with the aid of the Tagger technique. **Conclusion:** It could be concluded that a combination of knowledge about anatomy, techniques and substances yield satisfactory results in 3D filling and hermetic seal of root canals, which leads to successful endodontic treatment.

Keywords: Transverse anatomy. Endodontics. Root canal filling.

How to cite this article: Marion JJC, Batista DM, Murata SS, Souza V, Holland R. Factors that favor tridimensional and hermetic obturation of root canal system. *Dental Press Endod.* 2014 Jan-Apr;4(1):26-33. doi: <http://dx.doi.org/10.14436/2178-3713.V4.N1.026-033.oar>.

¹PhD resident in Clinical Dentistry/Endodontics, State University of Campinas (UNICAMP). Professor, Department of Endodontics, Brazilian Dental Association (ABO) and Ingá College (UNINGÁ)

²Degree in Dentistry, Ingá College (UNINGÁ).

³PhD in Pediatric Dentistry, School of Dentistry – UNESP / Araçatuba.

⁴Professor of Endodontics, Department of Restorative Dentistry, School of Dentistry – UNESP / Araçatuba.

» Patients displayed in this article previously approved the use of their facial and intraoral photographs.

» The authors report no commercial, proprietary or financial interest in the products or companies described in this article.

Submitted: November 01, 2013. Revised and accepted: November 08, 2013.

Contact address: Jefferson J. C. Marion
E-mail: jefferson@jmarion.com.br

Introduction

The main purposes of endodontic therapy are root canal cleaning achieved by removing pulp tissue; and root canal disinfection achieved by removing remaining necrotic pulp tissue, microorganisms and infected dentin.^{1,2} Knowing the inner anatomy of root canals is key for endodontic treatment success. Such knowledge allows endodontic treatment planning; however, anatomical changes may significantly hinder successful endodontic results.³ Despite being extensively studied, the clinical relevance of tooth anatomy has to be highlighted so as to emphasize its potential influence over the final results of endodontic treatment.⁴ According to Guldener,⁵ one must remember that a tooth root has one or more than one main canals which may be divided into several accessory root canals.

Knowledge about tooth anatomy is of paramount importance for the dental specialties. In other words, knowledge of canal morphology and its frequent variations before satisfactory infection control and advanced operating techniques strongly influence daily dental practice.

According to Pucci and Reig,⁶ the root portion of the pulp cavity often presents furcations or fusions. Depending on their location, root furcations are named as: main canal, collateral canal, lateral canal, secondary canal, accessory canal, intercanal, recurrent canal, reticular canals or apical deltas. Moreover, the classification by Kuttler⁷ also mentions an accessory canal in the furcation. According to De Deus,⁸ it is important to highlight, especially to beginners in the endodontic practice, that the aforementioned furcations are not found in one single root at the same time.

Duque⁹ assessed 15 endodontically treated teeth using a viewing box with a magnifying glass, and found no lateral canal. Nevertheless, De Deus,¹⁰ Vertucci¹¹ and Hess¹² proved that the aforementioned furcations are observed in all types of teeth. However, for Weine¹³ they are not enough demonstrated.

Rubach and Mitchell¹⁴ revealed that lateral and accessory canals are found in 45% of teeth with periodontal lesions, regardless of disease severity. Likewise, Kirkham¹⁵ concluded that lateral canals were found in 23% of periodontally compromised teeth. The author also highlighted that these pathways favor bacterial accumulation and may lead to pulp contamination, both of which hinder periodontal healing.

According to Murata,¹⁶ the mechanical action of files during biomechanical preparation of root canal dentin walls not only increases light inside the root canal, but also causes dentin debris which may be associated with organic debris and, for this reason, need to be removed by irrigating solution and suction.

Silva et al¹⁷ assert that effective removal of organic and inorganic compounds of dentin debris (smear layer) relies on the use of EDTA followed by sodium hypochlorite. The use of EDTA to remove smear layer has been effectively demonstrated by the literature.¹⁸⁻²⁷

According to Panzarini et al,²⁸ removing the smear layer after root canal biomechanical preparation has four major advantages: enhances dentin permeability^{29,30} and, as a result, favors the deep action of intracanal dressing inside the dentin tubules and furcations of the main canal;³¹ 2) favors adhesion of the filling cement to the root canal dentin walls;³² 3) provides better marginal, apical sealing;^{33,34} 4) allows the filling cement to drain through the furcations of the main canal.^{35,36}

The study conducted by Fachin and Salles³⁷ revealed that cleaner dentin walls, as a result of removing the smear layer, allow better adhesion of the filling cement to dentin tubules, which results in a better sealing.

As for Spångberg,³⁸ since the oral cavity is home for a wide variety of pathogenic microorganisms, the filling material must have physicochemical properties that favor permanent and effective root canal sealing, and, as a consequence, protects the apical and periapical tissues against microbial contamination and further infection and inflammation.

Some authors^{39,40,41} lay the responsibility of unsuccessful endodontic therapy to incomplete or inappropriate root canal filling.

With a view to increasing adhesion of filling material to root canal system walls, endodontic cements have also been used.⁴²

Whenever thermoplastic filling is carried out, we consciously decrease the amount of cement.⁴³ Nevertheless, the thin pellicle layer of cement must be well adhered to the dentin walls and, if possible, must have entered into the dentin tubules.⁴⁴ According to Schilder⁴⁵ as well as Sen, Piskin and Baran,⁴² the filling cement is the fragile portion of filling procedures; however, it plays an important role in controlling apical percolation as it flows through root

canal furcations and, as a result, provides the irregularities of the dentin-filling material interface with better sealing.

The literature describes that there is a tendency for future cementing material to adhere to dentin tubules, fuse with organic and inorganic dentin, destroy or neutralize microorganisms as well as their by-products, induce cementum neof ormation and strengthen the root canal system. However, it is worth noting that, from this point of view, the endodontic cements used nowadays are considered inappropriate.⁴⁶

We believe that the main canal must be well prepared and the root canal system must be properly and effectively filled. Based on that, the objective of this study is to demonstrate the possibility and importance of filling different types of accessory canals of different teeth. We report 11 cases in which different accessory canals at different locations were filled with 1% sodium hypochlorite and 17% trisodium EDTA irrigation, as well as three types of endodontic cement that, all together, favored tridimensional and hermetic sealing of the root canal system.

Methods and case report

This study comprises a sample of endodontic cases selected to demonstrate the possibility and need for tridimensional and hermetic filling of the root canal system. Biomechanical preparation and filling techniques were the same, whereas the endodontic cement varied for all cases.

Root canals were accessed by coronal portion. Tooth length was determined by odontometry. Working length was established at 1 mm below the radiographic root apex. Root canals underwent instrumentation carried out by means of the crown-down technique. After instrumentation, a FlexoFile file #15 (Dentsply Indústria e Comercio Ltda, Petrópolis/RJ - Brazil) was inserted into the apical foramen with a view to cleaning the cemental canal. Irrigating solution with 1% sodium hypochlorite (ASFER Indústria Química Ltda, São Caetano do Sul/SP - Brazil) was used as an additional instrumentation agent during root canal biomechanical preparation. After biomechanical preparation, root canals were dried with absorbent paper points (Dentsply) and filled with 17% trisodium EDTA (Biodinâmica Quim. e Farm. Ltda, Ibiporã/PR - Brazil) for 3 minutes, with mechanical

agitation to remove the smear layer.⁴⁷ A last irrigation procedure was carried out with saline solution (Eurofarma Laboratórios Ltda, São Paulo/SP - Brazil), after which root canals were dried with absorbent paper points and filled with intracanal dressing. Before final root canal filling, intracanal dressing was removed and the protocol with EDTA was repeated.

Root canal filling was performed by means of the Tagger's hybrid technique (first introduced in 1984). This technique associates active lateral condensation of the apical third to McSpadden thermo-mechanical compaction in the cervical and middle thirds, with vertical condensation carried out by Paivaconductors (GolgranIndústria e Comércio de Instrumentos Odontológicos, São Paulo/SP - Brazil). The aforementioned procedures allow sealing of the root canal system to be performed as best as possible. The following filling cements were used: zinc oxide (Biodinâmica Quim. e Farm. Ltda, Ibiporã/PR - Brazil) (Fig 1), Endométhasone (Septodont, France) (Fig 2) and Sealapex (SybronEndo [SBS] Glendora, USA) associated with iodoform (Quimidrol Com. Ind. Imp. Ltda) (Fig 3). After the filling procedure, all teeth were radiographed. Radiographic exams revealed main canal and its furcations completely filled with filling cement.

Discussion

The literature does not reach a consensus regarding the need for routine filling of lateral and accessory canals. In spite of that, detailed knowledge of the inner anatomical shape of root canal systems, as well as of instrumentation, irrigation and filling techniques, in addition to the use of biologically compatible material are key to the success of endodontic treatment. With regard to the inner anatomical shape of root canal systems, several researches⁴⁸⁻⁵¹ have highlighted the high incidence of furcations in the main canal. De Deus¹⁰ observed that 27.4% of the teeth studied had some type of furcation which were generally found in the apical region of the root.

According to Hatton et al,⁵² furcations of the main canal do not usually affect bone repair after treatment. Conversely, the study by Nicholls⁵³ reveals that periapical lesions of the main canal occurred in 4.4% of teeth studied. The cases report herein corroborate the study by Nicholls,⁵³ since periapical tissue repair was observed in cases presenting periapical lesion (Figs 1A, 2A, 2B, 2D, 3A, 3C, 3D).




Figure 1. A) Main canal (the most important and complex root canal, it goes through the long tooth axis and may directly reach the root apex) and apical delta filling (multiple furcations of the main canal in the apical region, they originate multiple foramina that replace the main foramen) with zinc oxide filling cement and Eugenol. **B)** Main canal and recurrent canal filling carried out with zinc oxide filling cement and Eugenol. The recurrent canal originates in the main canal, follows a long trajectory in the dentin and returns to the main canal before reaching the root apex).




Figure 2. A) Main canal and lateral canal filling carried out with Endométhasone filling cement. The lateral canal originates in the main canal and reaches the outer surface of the root in its cervical or middle thirds). **B)** Main canal and secondary canal filling carried out with Endométhasone filling cement. The secondary canal originates in the apical portion of the main canal and directly reaches the outer surface of the root). **C)** Main canal and secondary canal filling with Endométhasone filling cement. **D)** Main canal and lateral canal filling with Endométhasone filling cement.




Figure 3. **A)** Main canal and lateral canal filling with Sealapex filling cement. **B)** Main canal apical delta, secondary canal and accessory canal filling carried out with Sealapex filling cement. The accessory canal originates in a secondary canal and reaches the outer surface of the apical cementum. **C)** Main canal and secondary canal filling with Sealapex filling cement. **D)** Main canal and secondary canal filling with Sealapex filling cement. **E)** Main canal and accessory canal in the furcation filling carried out with Sealapex filling cement. The accessory canal in the furcation originates on the floor of the pulp chamber and reaches root bi or trifurcation.

Wada et al⁵⁴ conducted a study in which 70% of cases had apical furcations of the main canal. They also observed the impossibility of mechanically accessing these furcations, highlighting the need for performing a few procedures that allow the intracanal dressing to effectively act and ease the flow of filling material throughout the entire root canal system. Studies by Goldman et al⁵⁵ reveal that, after instrumentation, root canal walls are covered by a residual layer known as smear layer. It consists of organic and inorganic matter and hinders intracanal medication and filling material penetration inside the dentin tubules and small furcations of the main canal. De Deus

et al⁵⁶ concluded that the smear layer negatively affects the penetration capacity of endodontic cement inside dentin tubules.

EDTA has proved more effective^{26,27} than 2.5% sodium hypochlorite in removing smear layer as well as cleaning dentin tubules and the opening of accessory canals. The study by Holland et al³⁶ demonstrates that radiographic exams of filled root canals of teeth with lateral and apical furcations reveal the presence of filling cement in 90% of cases when, before performing the filling procedure, EDTA was applied for 3 minutes inside the root canal. On the other hand, the presence of filling cement occurred in 35% of

cases in which EDTA was not used. In the present study, irrigation with sodium hypochlorite associated with EDTA applied for 3 minutes increased the activity of filling cement in a higher number of cases as well as in accessory canals after the filling procedure, thus corroborating Holland et al.³⁶

Tam and Yu⁵⁰ assert that failure to seal accessory canal openings may lead to unsuccessful treatment results. Guimarães et al.⁵⁸ concluded that drainage of endodontic cement is responsible for closely fitting the material to the irregularities of the root canal walls, an essential factor that favors hermetic filling.

With a view to enhancing the properties of filling cements, some authors^{36,59,60} increased/decreased its ratio or added new substances. In the present study, root canals were filled with three different types of cement — Zinc oxide, Eugenol and Endométhasone — which had the following physical and chemical properties: drainage; working/setting time; and widely studied solubility and radiopacity in accordance with ISO-DIS 6876.2 (re-edition of ADA n° 57). Sealapex cement, however, was criticized with regard to some of its physical and chemical properties, including lower radiopacity in comparison to what is required by ADA n° 57. For this reason, iodoform was added to its original formulation.⁶¹ In the cases reported herein, Sealapex cement was associated with iodoform so as to achieve higher radiopacity, as suggested by Holland et al.³⁶

Today, however, such association is no longer necessary, given that the original formulation of Sealapex cement was modified with barium sulfate replaced by bismuth trioxide. According to Guimarães et al.,⁵⁸ drainage of Sealapex cement is nearly null, for this reason, this type of cement hardly enters into accessory canals. Nevertheless, Holland and Murata⁶² filled root canals of extracted human teeth with natural apical and artificial lateral furcations. Their study revealed that after filling performed by means of lateral condensation, the

cement drained into the furcations of all specimens. The aforementioned data explain the results yielded in the present study, since the cases presented herein demonstrate that even though Sealapex cement has lower drainage properties in comparison to other types of cement, it has ideal consistency when properly proportioned and applied, which allows it to enter into the root canal system.

Finally, root canal system filling also depends on the appropriate use of the technique which eliminates the empty gaps of the main canal and, as a result, favors cement drainage. We have produced successful results by initially performing lateral condensation of the apical third, followed by thermoplastification of gutta-percha cones carried out by means of McSpadden condensers. Nevertheless, it is worth noting that using demineralizing substances, good drainage cement and good filling technique significantly increase the chances of performing successful filling procedures; however, they do not assure success in 100% of cases.³⁶

Conclusion

Based on the results of this study it is reasonable to conclude that:

- » Detailed knowledge of the inner anatomical shape of root canal systems is essential for clinicians who aim at performing successful endodontic treatment.
- » The main canal and its furcations may be effectively filled after decalcifying solutions are applied (EDTA).
- » The physical, chemical and biological properties of filling cements must be considered when carefully choosing and correctly applying them.
- » The filling technique of choice has to provide hermetic sealing of the main root canal. The latter has to allow filling cement to drain to the root canal system during lateral and vertical condensation associated with Tagger filling technique.

References

- Rizzardo A. Estudo dos efeitos de algumas soluções irrigadoras sobre a microdureza dentinária [dissertação]. Passo Fundo (RS): Universidade Passo Fundo; 2001. 47 p.
- Nunes E. Influência de técnicas de instrumentação na penetração de cânulas irrigadoras em canais radiculares [dissertação] Bauru (SP): Universidade de São Paulo; 1993.
- Constante IGT, Albergaria S, Moura AAM. Estudo comparativo das alterações anatômicas promovidas pelas técnicas do Preparo Progressivo, Escalonada e Seriada em canais méso-vestibulares de dentes molares inferiores com curvatura severa. *Rev Inst Ciênc Saúde*. 2006;24(3):207-14.
- Gesteira MFM, Saraiva MO, Silva SJA. Estudo da anatomia interna de pré-molares inferiores pelo método da diafanização. *J Bras Endod*. 2006;6(24):78-83.
- Guldener PHA. Endodontia: diagnóstico e tratamento. Barcelona: Springer-Verlag; 1995.
- Pucci FM, Reig R. Conductos radiculares: anatomía, patología y terapia. Montevideo: Ed. A. Barreiro y Ramos; 1945.
- Kuttler Y. Endodontia prática. México: Alpha; 1961.
- De Deus QD. Topografia da cavidade pulpar: contribuição ao seu estudo. Belo Horizonte: Doctorate Thesis; 1960.
- Duque TM. Análise microbiológica por PCR, quantificação de citocinas e monitoramento inflamatório em pacientes com doença periodontal crônica [dissertação]. Piracicaba (SP): Universidade Estadual de Campinas; 2012.
- De Deus QD. Frequency, location, and direction of the lateral, secondary, and accessory canals. *J Endod*. 1975;1(11):361-6.
- Vertucci FJ. Root canal anatomy of the human permanent teeth. *Oral Surg Oral Med Oral Pathol*. 1984;58(5):589-99.
- Hess W. The anatomy of the root canals of the teeth of the permanent dentition. New York: William Wood; 1925.
- Weine FS. Endodontic therapy. 4th ed. St. Louis: Mosby; 1989. p. 245-51.
- Ruback WC, Mitchell DF. Periodontal disease accessory canals and pulp pathosis. *J Periodontol*. 1965;36:34-8.
- Kirkham DB. The location and incidence of accessory pulpal canals in periodontal pockets. *J Am Dent Assoc*. 1975;91(2):353-6.
- Murata SS. Análise histomorfológica de dentes descuidados de case com rizogênese incompleta após biopulpectomia e obturação dos canais radiculares com hidróxido de cálcio em diferentes veículos [tese]. Araçatuba (SP): Universidade Estadual Paulista; 2006.
- Silva JM, Silveira A, Santos E, Prado L, Pessoa OF. Efficacy of sodium hypochlorite, ethylenediaminetetraacetic acid, citric acid and phosphoric acid in calcium hydroxide removal from the root canal: a microscopic cleanliness evaluation. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2011;112(6):820-4.
- McComb D, Smith DC. A preliminary scanning electron microscopic study of root canals after endodontic procedures. *J Endod*. 1975;1(7):238-42.
- McComb D, Smith DC, Beagrie GS. The results of in vivo endodontic chemomechanical instrumentation: a scanning electron microscopic study. *J Br Endod Soc*. 1976;9(1):11-8.
- Goldberg F, Abramovich A. Analysis of the effects of EDTA on the dentinal walls of the root canal. *J Endod*. 1977;3(3):101-5.
- Yamada RS, Armas A, Goldman M, Lin PS. A scanning electron microscopic comparison of a high volume final flush with several irrigating solutions: part 3. *J Endod*. 1983;9(4):137-42.
- Cameron JA. The use of ultrasonics in the removal of the Smear layer: a scanning electron microscope study. *J Endod*. 1983;9(7):289-92.
- Baumgartner JC, Marder CL. A scanning electron microscopic valuation of four root canal irrigation regimens. *J Endod*. 1987;13(4):147-57.
- Meryon SD, Tobias RS, Jakeman KJ. Smear removal agents: a quantitative study in vivo and in vitro. *J Prosthet Dent*. 1987;57(2):174-9.
- Cengiz T, Aktener BO, Piskin B. The effect of dentinal tubule orientation on the removal of Smear layer by root canal irrigant. A scanning electron microscopic study. *Int Endod J*. 1990;23(3):163-71.
- Onoda HK, Yoshinari GH, Pereira KF, Guerisoli DMZ. Estudo dos efeitos de algumas soluções irrigadoras sobre a microdureza dentinária. In: 10º Simposio Brasil-Japão; 2010 Out. 8-12. Campo Grande: Associação Brasil-Japão de Pesquisadores; 2010.
- Adigüzel O, Yiğit-Özer S, Kaya S, Uysal İ, Ganidağlı-Ayaz S, Akkuş Z. Effectiveness of ethylenediaminetetraacetic acid (EDTA) and MTAD on debris and smear layer removal using a self-adjusting file. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2011;112(6):803-8.
- Panzarini SR, Souza V, Holland R, Dezan Júnior E. Tratamento de dentes com lesão periapical crônica: influência de diferentes tipos de curativo de demora e do material obturador de canal radicular. *Rev Odontol UNESP*. 1998;27(2):509-26.
- Hampson EL, Atkinson AM. The relation between drugs used in root canal therapy and the permeability of the dentine. *Br Dent J*. 1964;116(12): 546-50.
- Zina O, Souza V, Holland R, Saliba O. Influence of chelant agents on dentin permeability and on the healing process of periapical tissues after root canal treatment. *Rev Odontol UNESP*. 1981;10(1-2):27-33.
- Holland R, Okabe JN, Souza V, Saliba O. Diffusion of corticosteroid-antibiotic solutions through human dentine. *Rev Odontol UNESP*. 1991;20(1):17-23.
- Okşan T, Aktener BO, Sen BH, Tezel H. The penetration of root canal sealers into dentinal tubules. A scanning electron microscopic study. *Int Endod J*. 1993;26(5):301-5.
- Holland R, Paula EC, Pereira ALS, Souza V, Saliba O. Infiltração marginal dos cimentos endodônticos. *RGD: Rev Gaúch Odontol*. 1991;39(6):413-6.
- Saunders WP, Saunders EM. The effect of smear layer upon the coronal leakage of gutta-percha root fillings and a glass ionomer sealer. *Int Endod J*. 1992;25(5):245-9.
- Goldberg F, Spielberg C. The effect of EDTAC and the variation of its working time analyzed with scanning electron microscopy. *Oral Surg Oral Med Oral Pathol*. 1982;53(1):74-7.
- Holland R, Silva ACF, Bazaglia AM, Barros VCL, Magro VM. Influência do uso de soluções descalcificadoras na obturação do sistema de canais radiculares. *Rev Bras Odontol*. 1988;155(2):16-22.
- Fachin EVF, Salles AA. Influência do tratamento com EDTA no embrocamento de dois cimentos resinosos às paredes dos canais radiculares. *Rev Facul Odontol Porto Alegre*. 2003;44(2):39-42.
- Spångberg LSW. Tratamento endodôntico de dentes sem periodontite apical. In: Orstavik D, Pitt Ford TR. Fundamentos da endodontia: prevenção e tratamento da periodontite apical. São Paulo: Ed. Santos; 2004. p. 211-41.
- Ingle JI. Root canal obturation. *J Am Dent Assoc*. 1956;53(1):47-55.
- Swartz DB, Skidmore AE, Griffin JA. Twenty years of endodontic success and failure. *J Endod*. 1983;9(5):198-202.
- Ngruyen NT. Obturation of the root canal system. In: Cohen S, Burns R. Pathways of the pulp. 6th ed. St. Louis: Mosby; 1994.
- Sen BH, Piskin B, Baran N. The effect of tubular penetration of root canal sealers on dye microleakage. *Int Endod J*. 1996;29(1):23-8.
- Yared GM, Bou-Dagher F. Sealing ability of the vertical condensation with different root canal sealers. *J Endod*. 1996;22(1):6-8.
- Kouvas V, Liolios E, Vassiliadis L, Parisiss-Messimeris S, Boutsioukis A. Influence of smear layer on depth of penetration of three endodontic sealers: a SEM study. *Endod Dent Traumatol*. 1998;14(4):191-5.
- Schilder H. Filling root canals in three dimensions. *Dent Clin North Am*. 1967;11:723-44.

46. Gutmann JL, Whitherspoon DE. Obturation of the cleaned and shaped root canal system. In: Cohen S, Burns RC. *Pathways of the Pulp*. 7th ed. St. Louis: Mosby; 1998. p. 258-361.
47. Holland R, Silva ACF, Bazaglia AM, Barros VCL, Magro VM. Influência do uso de soluções descalcificadoras na obturação do sistema de canais radiculares. *Rev Bras Odontol*. 1988;45(2):16-22.
48. Seltzer S. *Endodontology. Biologic considerations in endodontic procedures*. New York: McGraw Hill Book Company; 1971.
49. Hess W, Keller O. *Le tavole anatomiche*. Saronno: Edizioni scientifiche Oral B; 1988.
50. Tam A, Yu DC. Location of canal isthmus and accessory canals in the mesiobuccal root of maxillary first permanent molars. *J Can Dent Assoc*. 2002;68(1):28-33.
51. Martos J, Lubian C, Silveira LF, Suita de Castro LA, Ferrer Luque CM. Morphologic analysis of the root apex in human teeth. *J Endod*. 2010;36(4):664-7.
52. Hatton EH, Skillen WG, Moen OH. Histologic findings in teeth with treated and filled root canals. *J Am Dent Assoc*. 1928;15:56.
53. Nicholls E. Lateral radicular disease due to lateral branching of the root canal. *Oral Surg Oral Med Oral Pathol*. 1963;16:839-45.
54. Wada M, Takase T, Nakanuma K, Arisue F, Nagahana F, Yamazaki M. Clinical study of refractory apical periodontitis treated by apicectomy. Part I: root canal morphology of resected apex. *Int Endod J*. 1998;31(1):51-6.
55. Goldman LB, Goldman M, Kronman JH, Yeck SL. The efficacy of several endodontic irrigating: a scanning electron microscopic study. *Oral Surg*. 1981;52:197-204.
56. De Deus G, Gurgel Filho ED, Ferreira CM, Coutinho Filho T. Penetração intratubular de cimentos endodônticos. *Pesqui Odontol Bras*. 2002;16(4):332-6.
57. Goldberg F, Massone JE, Spielberg C. Effect of irrigation solutions on the filling of lateral root canals. *Endod Dent Traumatol*. 1986;2(2):65-6.
58. Guimarães CCP, Sampaio JMP, Sato EFL, Collesi RR. Avaliação do escoamento de seis cimentos endodônticos. *Rev Odontol Univ St Amaro*. 1999;4(1):4-10.
59. Pécora JD, Ribeiro RG, Guerisoli DMZ, Brabizam JU. Influência da espatulação de dois cimentos à base de OZE na obturação de canais laterais. *Pesq Odont Bras*. 2002;16(2):127-30.
60. Duarte MAH. *Avaliação de algumas propriedades físico-químicas do cimento AH Plus puro e acrescido de hidróxido de cálcio [tese]*. Bauru (SP): Universidade de São Paulo; 1999.
61. Almeida JF, Gomes BP, Ferraz CC, Souza-Filho FJ, Zaia AA. Filling of artificial lateral canals and microleakage and flow of five endodontic sealers. *Int Endod J*. 2007;40(9):692-9.
62. Holland R, Murata SS. Obturação de canais radiculares com cimentos à base de hidróxido de cálcio. *Rev APCD*. 1995;49(3):221-4.